

Five Things Physicians and Patients Should Question

1 **Avoid routine multiple daily self-glucose monitoring in adults with stable type 2 diabetes on agents that do not cause hypoglycemia.**

Once target control is achieved and the results of self-monitoring become quite predictable, there is little gained in most individuals from repeatedly confirming. There are many exceptions, such as for acute illness, when new medications are added, when weight fluctuates significantly, when A1c targets drift off course and in individuals who need monitoring to maintain targets. Self-monitoring is beneficial as long as one is learning and adjusting therapy based on the result of the monitoring.

2 **Don't routinely measure 1,25-dihydroxyvitamin D unless the patient has hypercalcemia or decreased kidney function.**

Many practitioners become confused when ordering a vitamin D test. Because 1,25-dihydroxyvitamin D is the active form of vitamin D, many practitioners think that measuring 1,25-dihydroxyvitamin D is an accurate means to estimate vitamin D stores and test for vitamin D deficiency, which is incorrect. Current Endocrine Society guidelines recommend screening for vitamin D deficiency in individuals at risk for deficiency.

Serum levels of 1,25-dihydroxyvitamin D have little or no relationship to vitamin D stores but rather are regulated primarily by parathyroid hormone levels, which in turn are regulated by calcium and/or vitamin D. In vitamin D deficiency, 1,25-dihydroxyvitamin D levels go up, not down.

Unregulated production of 1,25-dihydroxyvitamin D (i.e., sarcoidosis, granulomatous diseases) is an uncommon cause of hypercalcemia; this should be suspected if blood calcium levels are high and parathyroid hormone levels are low and confirmed by measurement of 1,25-dihydroxyvitamin D. The enzyme that activates vitamin D is produced in the kidney, so blood levels of 1,25-dihydroxyvitamin D are sometimes of interest in patients on dialysis or with end-stage kidney disease. There are few other circumstances, if any, where 1,25-dihydroxyvitamin D testing would be helpful.

Serum 25-hydroxyvitamin D levels may be overused, but when trying to assess vitamin D stores or diagnose vitamin D deficiency (or toxicity), 25-hydroxyvitamin D is the correct test.

3 **Don't routinely order a thyroid ultrasound in patients with abnormal thyroid function tests if there is no palpable abnormality of the thyroid gland.**

Thyroid ultrasound is used to identify and characterize thyroid nodules, and is not part of the routine evaluation of abnormal thyroid function tests (over- or underactive thyroid function) unless the patient also has a large goiter or a lumpy thyroid. Incidentally discovered thyroid nodules are common. Overzealous use of ultrasound will frequently identify nodules, which are unrelated to the abnormal thyroid function, and may divert the clinical evaluation to assess the nodules, rather than the thyroid dysfunction. Imaging may be needed in thyrotoxic patients; when needed, a thyroid scan, not an ultrasound, is used to assess the etiology of the thyrotoxicosis and the possibility of focal autonomy in a thyroid nodule.

4 **Don't order a total or free T3 level when assessing levothyroxine (T4) dose in hypothyroid patients.**

T4 is converted into T3 at the cellular level in virtually all organs. Intracellular T3 levels regulate pituitary secretion and blood levels of TSH, as well as the effects of thyroid hormone in multiple organs; a normal TSH indicates an adequate T4 dose. Conversion of T4 to T3 at the cellular level may not be reflected in the T3 level in the blood. Compared to patients with intact thyroid glands, patients taking T4 may have higher blood T4 and lower blood T3 levels. Thus the blood level of total or free T3 may be misleading (low normal or slightly low); in most patients a normal TSH indicates a correct dose of T4.

5 **Don't prescribe testosterone therapy unless there is biochemical evidence of testosterone deficiency.**

Many of the symptoms attributed to male hypogonadism are commonly seen in normal male aging or in the presence of comorbid conditions. Testosterone therapy has the potential for serious side effects and represents a significant expense. It is therefore important to confirm the clinical suspicion of hypogonadism with biochemical testing. Current guidelines recommend the use of a total testosterone level obtained in the morning. A low level should be confirmed on a different day, again measuring the total testosterone. In some situations, a free or bioavailable testosterone may be of additional value.

How This List Was Created

Members of The Endocrine Society (Society) along with representatives of the American Association of Clinical Endocrinologists (AACE) formed a joint task force to identify tests or procedures which should only be used in specific circumstances. The task force identified several items for possible inclusion. Subsequent discussions compared the evidence supporting each item, the value of the recommendation to practitioners and the potential for cost savings. Members of the Society's Clinical Affairs Core Committee and AACE leadership also reviewed the initial list. Using the above criteria, the task force voted for their top five recommendations from the original list. The Society's Council and AACE's Board of Directors approved the final list for submission to the *Choosing Wisely*[®] campaign.

The Endocrine Society disclosure and conflict of interest policies can be found at www.endocrine.org.

*The American Association of Clinical Endocrinologists withdrew from the *Choosing Wisely*[®] campaign on May 26, 2015.

Sources

- Davidson MB, Castellanos M, Kain D, Duran P. The effect of self monitoring of blood glucose concentrations on glycated hemoglobin levels in diabetic patients not taking insulin: a blinded, randomized trial. *Am J Med.* 2005;118:422–5.
Farmer A, Wade A, Goyder E, Yudkin P, French D, Craven A, Holman Rury, Kinmonth AL, Neil A. Impact of self monitoring of blood glucose in the management of patients with non-insulin treated diabetes: open parallel group randomized trial. *BMJ.* 2007;335:132–40.
O'Kane MJ, Bunting B, Copeland M, Coates VE; ESMON study group. Efficacy of self monitoring of blood glucose in patients with newly diagnosed type 2 diabetes (ESMON study): randomized controlled trial. *BMJ.* 2008;336:1174–7.
- Bikle D, Adams J, Christakos S. Primer on the metabolic bone diseases and disorders of mineral metabolism. Washington: American Society for Bone and Mineral Research. c2008. Chapter 28, Vitamin D: production, metabolism, mechanism of action, and clinical requirements. P. 141–9.
Holick MF. Vitamin D deficiency. *N Engl J Med.* 2007;357:266–81.
Holick MF, Binkley NC, Bischoff-Ferrari HA, Gordon CM, Hanley DA, Heaney RP, Murad MH, Weaver CM; Endocrine Society. Evaluation, treatment, and prevention of vitamin D deficiency: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab.* 2011 Jul;96(7):1911–30.
- Bahn RS, Burch HB, Cooper DS, Garber JR, Greenlee MC, Klein I, Laurberg P, McDougall IR, Montori VM, Rivkees SA, Ross DS, Sosa JA, Stan MN; American Thyroid Association; American Association of Clinical Endocrinologists. Hyperthyroidism and other causes of thyrotoxicosis: management guidelines of the American Thyroid Association and American Association of Clinical Endocrinologists. *Thyroid.* 2011;21:593–646.
Garber JR, Cobin RH, Gharib H, Hennessey JV, Klein I, Mechanick JI, Pessah-Pollack R, Singer PA, Woeber KA. Clinical practice guidelines for hypothyroidism in adults: cosponsored by the American Association of Clinical Endocrinologists and the American Thyroid Association. *Endocr Pract.* 2012; Sep 11:1–207.
- Garber JR, Cobin RH, Gharib H, Hennessey JV, Klein I, Mechanick JI, Pessah-Pollack R, Singer PA, Woeber KA. Clinical practice guidelines for hypothyroidism in adults: cosponsored by the American Association of Clinical Endocrinologists and the American Thyroid Association. *Endocr Pract.* 2012; Sep 11:1–207.
- Bhasin S, Cunningham GR, Hayes FJ, Matsumoto AM, Snyder PJ, Swerdloff RS, Montori VM. Testosterone therapy in adult men with androgen deficiency syndromes: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab.* 2006 Jun;91(6):1995–2010.
Wu FCW, Tajar A, Beynon JM, Pye SR, Silman AJ, Finn JD, O'Neill TW, Bartfai G, Casanueva FF, Forti G, Giwercman A, Han TS, Kula K, Lean ME, Pendleton N, Punab M, Boonen S, Vanderschueren D, Labrie F, Huhtaniemi IT; EMAS Group. Identification of late-onset hypogonadism in middle-aged and elderly men. *N Engl J Med.* 2010 Jul 8;363(2):123–35.

About the ABIM Foundation

The mission of the ABIM Foundation is to advance medical professionalism to improve the health care system. We achieve this by collaborating with physicians and physician leaders, medical trainees, health care delivery systems, payers, policymakers, consumer organizations and patients to foster a shared understanding of professionalism and how they can adopt the tenets of professionalism in practice.

To learn more about the ABIM Foundation, visit www.abimfoundation.org.



About The Endocrine Society

- Founded in 1916, The Endocrine Society is the world's oldest, largest, and most active organization devoted to research on hormones and the clinical practice of endocrinology. The Society is an international body with more than 16,000 members from over 100 countries, and represents the full range of disciplines associated with endocrinologists: clinicians, researchers, educators, fellows and students, industry professionals and health professionals who are involved in the field of endocrinology. Our members are dedicated to the research and treatment of the full range of endocrine disorders: diabetes, reproduction, infertility, osteoporosis, thyroid disease, obesity/lipids, growth hormone, pituitary tumors and adrenal insufficiency.

Visit The Endocrine Society at www.endocrine.org.



For more information or to see other lists of Five Things Physicians and Patients Should Question, visit www.choosingwisely.org.